arXiv:1609.04771v1 [math.CA] 4 Sep 2016

Formula of Volume of Revolution with Integration by Parts and Extension

Yi Liu , Jingwei Liu

School of Mathematics and Statistics, Beijing Institute of Technology ,Beijing 100081,P.R China
Email : liuy0605@126.com.

School of Mathematics and System Sciences, Beihang University, Beijing, 100083,P.R China
Email : liujingwei03@tsinghua.org.cn.

Abstract. A calculation formula of volume of revolution with integration by parts of definite integral is derived based on monotone function, and extended to a general case that curved trapezoids is determined by continuous, piecewise strictly monotone and differential function. And, two examples are given, ones curvilinear trapezoids is determined by Kepler equation, and the other curvilinear trapezoids is a function transmuted from Kepler equation.

In [1] (p494,ex71), if $f(x)$ is one-to-one and therefore has an inverse function $x = g(y)$, where $x \in [a, b]$,$a > 0$ and $y \in [c, d]$. Let S be the solid obtained by rotating about the y-axis the region bounded by $y = f(x)$ [where $f(x) \geq 0$], $y = 0$, $x = a$ and $x = b$ (which is also called curvilinear trapezoid [2]). Using cylindrical shells and slicing method, the volume of the given solid is

$$V = \int_a^b 2\pi x f(x)dx = \pi b^2 d - \pi a^2 c - \int_c^d \pi[g(y)]^2 dy \tag{1}$$

This formula can also be proved by integration by parts with the increasing case of f. However, there is slight divarication in formula (1).

i) If we guarantee the condition $y \in [c, d]$, formula (1) does not hold in case of decreasing f, as under this condition $c = f(b)$, $d = f(a)$.
ii) If we ensure that the condition $c = f(a)$, $d = f(b)$ is established regardless of increasing or decreasing of f, the condition $y \in [c, d]$ does not hold in case of decreasing of f, as $c > d$. However, formula (1) still holds in mathematical style.

1. VOLUME OF REVOLUTION AND EXTENSION. A reasonable expression would be as follows.

Theorem 1. *Let $y = f(x)$ be nonnegative, continuous, differential and strictly monotone on [a,b] ($a > 0$), it's inverse function is $x = g(y)$, $y \in [c, d]$. Then,*

$$\pi \int_c^d [g(y)]^2 dy = sgn(f(b) - f(a)) \left\{ \pi[b^2 f(b) - a^2 f(a)] - 2\pi \int_a^b x f(x)dx \right\} \tag{2}$$

$$\pi \int_a^b [f(x)]^2 dx = sgn(g(d) - g(c)) \left\{ \pi[d^2 g(d) - c^2 g(c)] - 2\pi \int_c^d y g(y)dy \right\} \tag{3}$$

where $y = sgn(x)$ is the sign function.

Proof. (I) Let $y = f(x)$ be strictly monotone increasing(fig1),then $c = f(a)$, $d = f(b)$, $c < d$, and

$$\begin{aligned} V_Y &= \pi \int_c^d [g(y)]^2 dy = \pi \int_a^b x^2 df(x) = \pi \left[x^2 f(x)|_a^b - \int_a^b f(x) dx^2 \right] \\ &= \pi \left[b^2 f(b) - a^2 f(a) \right] - 2\pi \int_a^b x f(x) dx \end{aligned} \tag{4}$$



Figure 1. $y = f(x)$ is strictly monotone increasing. **Figure 2.** $y = f(x)$ is strictly monotone decreasing.

(II) Let $y = f(x)$ be strictly monotone decreasing (fig 2), then $d = f(a)$, $c = f(b)$, ($c < d$), and

$$\begin{aligned} V_Y &= \pi \int_c^d [g(y)]^2 dy = \pi \int_b^a x^2 df(x) = -\pi \int_a^b x^2 df(x) \\ &= -\pi \left[x^2 f(x)|_a^b - \int_a^b f(x) dx^2 \right] \\ &= 2\pi \int_a^b x f(x) dx - \pi \left[b^2 f(b) - a^2 f(a) \right] \end{aligned} \tag{5}$$

Similarly, we can prove the other two cases of volume of solids of revolution with integration by parts as follows.

(III) If $x = g(y)$ is strictly monotone increasing (fig 1), then $a = g(c)$, $b = g(d)$, ($a < b$), and

$$\begin{aligned} V_X &= \pi \int_a^b [f(x)]^2 dx = \pi \int_c^d y^2 dg(y) = \pi \left[y^2 g(y)|_c^d - \int_c^d g(y) dy^2 \right] \\ &= \pi \left[d^2 g(d) - c^2 g(c) \right] - 2\pi \int_c^d y g(y) dy \end{aligned} \tag{6}$$

(IV) If $x = g(y)$ is strictly monotone decreasing (fig 2), then $b = g(c)$, $a = g(d)$,

($a < b$), and

$$\begin{aligned} V_X &= \pi \int_a^b [f(x)]^2 dx = \pi \int_d^c y^2 dg(y) = -\pi \int_c^d y^2 dg(y) \\ &= -\pi \left[y^2 g(y)|_c^d - \int_c^d g(y) dy^2 \right] \\ &= 2\pi \int_c^d yg(y) dy - \pi \left[d^2 g(d) - c^2 g(c) \right] \end{aligned} \tag{7}$$

Hence, this ends the proof. ■

If $y = f(x)$ is continuous and not strictly monotone on $[a, b]$, and $[a, b]$ can be divided into several strictly monotone intervals of $f(x)$, we have the following lemma.

Lemma 1. Suppose that $y = f(x)$is continuous and piecewise strictly monotone on $[a, b]$, if $f(a) \neq f(b)$, $\min\{f(a), f(b)\} < f(x) < \max\{f(a), f(b)\}$, $x \in (a, b)$, and $f(x)$ is not strictly monotone on $[a, b]$, the number of local extreme value points in (a, b) is even.

Proof. We prove the lemma in two cases, $f(a) < f(b)$, and $f(a) > f(b)$.

In case of $f(a) < f(b)$, since $f(x) > f(a)$, $x \in (a, b)$, the nearest extreme value point to $x = a$ is a local maximum value point. Since $f(x) < f(b)$, $x \in (a, b)$, the nearest extreme value point to $x = b$ is a local minimum value point. As $y = f(x)$ is continuous and piecewise strictly monotone on $[a, b]$, the number of extreme value points should be even. OR, using method of reduction to absurdity, if the number of extreme values was odd, the last extreme value point (nearest extreme value point to $x = b$) might be a local maximum extreme value point, it is a contradiction.

Similarly, we can prove the other case. Hence, we complete the proof. ■

Then, the volume of curvilinear trapezoidal composed by curved edge $y = f(x)$ in term of y-axis, and rotating with y-axis can reach the following conclusion.

Theorem 2. *Let $y = f(x)$ be continuous, piecewise strictly monotone and differential on $[a, b]$. $\Gamma$is the curve determined by $y = f(x)$ on $[a, b]$. Denote $c = \min(f(a), f(b))$, $d = \max(f(a), f(b))$. Suppose that both $y = c$ and $y = d$ $(c < d)$ intersect $y = f(x)$ in only one point on $[a, b]$ respectively. Then the volume of the plane graph along $x = 0$, $y = c$, Γ, $y = d$ rotating with y-axis is*

$$V_Y = sgn(f(b) - f(a)) \left\{ \pi \left[b^2 f(b) - a^2 f(a) \right] - 2\pi \int_a^b xf(x) dx \right\} \tag{8}$$

Proof. According to Lemma 1, since $y = f(x)$ is continuous, and strictly piecewise monotone on $[a, b]$, assume that the strictly piecewise monotone intervals of $y = f(x)$ are $[x_i, x_{i+1}]$, $i = 0, 1, \cdots, n$, $x_0 = a$, $x_{n+1} = b$, $i = 1, 2, \cdots, n$, and $[a, b] = \bigcup_{i=0}^{n} [x_i, x_{i+1}]$, where $x_1, \cdots, x_n$ are extreme value points of f.

(1) If $f(b) > f(a)$ (fig3), as both $y = c$ and $y = d$ $(c < d)$ intersect $y = f(x)$ only once on $[a, b]$ respectively, $y = f(x)$ is strictly increasing on both $[x_0, x_1]$ and $[x_n, x_{n+1}]$, where n is even (if $n = 0$, Theorem 2 degrades to Theorem 1) and $c = f(a)$, $d = f(b)$.

Denote $\Gamma_i : y = f(x), x \in [x_i, x_{i+1}]$, and V_{Γ_i} is the volume of curved trapezoid Γ_i with y -axis rotating with y-axis. Since $y = f(x)$ is strictly monotone and differential



Figure 3. f is continuous and piecewise strictly monotone with $f(b) > f(a)$.



Figure 4. f is continuous and piecewise strictly monotone with $f(b) < f(a)$.

on $[x_i, x_{i+1}]$, Theorem 1 would be employed in the volume of solids of revolution on each curvilinear trapezoidal determined by piecewise strictly monotone and differential curve. Then,

$$\begin{aligned} V_{\Gamma_{2k}} &= \pi[x_{2k+1}^2 f(x_{2k+1}) - x_{2k}^2 f(x_{2k})] - 2\pi \int_{x_{2k}}^{x_{2k+1}} x f(x) dx \\ V_{\Gamma_{2k+1}} &= 2\pi \int_{x_{2k+1}}^{x_{2k+2}} x f(x) dx - \pi[x_{2k+2}^2 f(x_{2k+2}) - x_{2k+1}^2 f(x_{2k+1})] \\ & \qquad\qquad k = 0, 1, \cdots, n/2. \end{aligned}$$

Using mathematical induction, we can obtain,

$$\begin{aligned} V_Y &= V_{\Gamma_0} - V_{\Gamma_1} + V_{\Gamma_2} + \cdots + (-1)^n V_{\Gamma_n} = \sum_{i=0}^{n} (-1)^{i-1} V_{\Gamma_i} \\ &= \sum_{i=0}^{n} \left\{ \pi[x_{i+1}^2 f(x_{i+1}) - x_i^2 f(x_i)] - 2\pi \int_{x_i}^{x_{i+1}} x f(x) dx \right\} \\ &= \pi[x_{n+1}^2 f(x_{n+1}) - x_0^2 f(x_0)] - 2\pi \int_{x_0}^{x_{n+1}} x f(x) dx \\ &= \pi[b^2 f(b) - a^2 f(a)] - 2\pi \int_a^b x f(x) dx \end{aligned} \tag{9}$$

(2) If $f(a) > f(b)$ (fig4), as both $y = c$ and $y = d$ $(c < d)$ intersect $y = f(x)$ one point on $[a, b]$ respectively. Then $y = f(x)$ is strictly decreasing on $[x_0, x_1]$ and $[x_n, x_{n+1}]$, where n is even, and $d = f(a)$, $c = f(b)$.

Denote $\Gamma_i : y = f(x), x \in [x_i, x_{i+1}]$,and V_{Γ_i} is the volume of curvilinear trapezoid of Γ_i in term of y-axis rotating with y-axis. Then,

$$\begin{aligned} V_{\Gamma_{2k}} &= 2\pi \int_{x_{2k}}^{x_{2k+1}} x f(x) dx - \pi[x_{2k+1}^2 f(x_{2k+1}) - x_{2k}^2 f(x_{2k})] \\ V_{\Gamma_{2k+1}} &= \pi[x_{2k+2}^2 f(x_{2k+2}) - x_{2k+1}^2 f(x_{2k+1})] - 2\pi \int_{x_{2k+1}}^{x_{2k+2}} x f(x) dx \\ & \qquad\qquad k = 0, 1, \cdots, n/2. \end{aligned}$$

Using mathematical induction, we can obtain,

$$\begin{aligned} V_Y &= V_{\Gamma_0} - V_{\Gamma_1} + V_{\Gamma_2} + \cdots + (-1)^n V_{\Gamma_n} = \sum_{i=0}^{n} (-1)^{i-1} V_{\Gamma_i} \\ &= \sum_{i=0}^{n} \left\{ 2\pi \int_{x_i}^{x_{i+1}} x f(x) dx - \pi [x_{i+1}^2 f(x_{i+1}) - x_i^2 f(x_i)] \right\} \\ &= 2\pi \int_{x_0}^{x_{n+1}} x f(x) dx - \pi [x_{n+1}^2 f(x_{n+1}) - x_0^2 f(x_0)] \\ &= 2\pi \int_a^b x f(x) dx - \pi [b^2 f(b) - a^2 f(a)] \end{aligned} \tag{10}$$

To sum up formulae(9)(10), the theorem is completely proved. ■

Similarly, we can obtain the following theorem:

Theorem 3. *Let $x = g(y)$ be continuous, piecewise strictly monotone and differential on $[c, d]$. Γ is the curve determined by $x = g(y)$ on $[c, d]$. Denote $a = \min(g(c), g(d))$, $b = \max(g(c), g(d))$. Suppose that both $x = a$ and $x = b$ $(a < b)$ intersect $x = g(y)$ in only one point on $[c, d]$ respectively. Then the volume of the plane graph along $y = 0$, $x = a$, Γ and $x = b$ rotating with x-axis is*

$$V_X = sgn(g(d) - g(c)) \left\{ \pi \left[d^2 g(d) - c^2 g(c) \right] - 2\pi \int_c^d y g(y) dy \right\} \tag{11}$$

2. APPLICATION. Example 1 For Kepler equation [3]: $y = x + \varepsilon \sin y$, $\varepsilon \in (0, 1)$, $y \in [0, 2\pi]$. (fig 5) (i) To calculate the volume of solids of revolution of curved trapezoid composed by Kepler curve, $y = 0, y = 2\pi$ and y-axis rotating with y-axis. (ii) To calculate the volume of solids of revolution of curved trapezoid composed by Kepler curve, $x = 0, x = 2\pi$ and x-axis rotating with x-axis.

Solution (i) For $x = g(y) = y - \varepsilon \sin y$,$\varepsilon \in (0, 1)$,

$$\begin{aligned} V_Y &= \pi \int_0^{2\pi} [g(y)]^2 dy = \pi \int_0^{2\pi} [y - \varepsilon \sin y]^2 dy \\ &= [\frac{y^3}{3} + 2\varepsilon(y \cos y - \sin y) + \varepsilon^2 (\frac{y}{2} - \frac{\sin 2y}{4})]|_0^{2\pi} \\ &= \frac{8}{3}\pi^3 + (4\varepsilon + \varepsilon^2)\pi \end{aligned}$$

(ii) Since $\dfrac{dx}{dy} = 1 - \varepsilon \cos y > 0$, the curve determined by Kepler function $x = g(y) = y - \varepsilon \sin y$ is strictly increasing in term of y on $[0, 2\pi]$. According to inverse function theorem of monotone continuous function, it's inverse function $y = f(x)$ exists in term of x on $[0, 2\pi]$ and is strictly increasing. Then, the function $y = f(x)$ determined by Kepler function is an implicit function and also transcendental equation.

Using formula(3), $a = 0$, $b = 2\pi$, $c = 0$, $d = 2\pi$, we obtain

$$\begin{aligned} V_X &= \pi \int_0^{2\pi} [f(x)]^2 dx = 8\pi^4 - 2\pi \int_0^{2\pi} yg(y)dy \\ &= 8\pi^4 - 2\pi \int_0^{2\pi} y(y - \varepsilon \sin y)dy \\ &= 8\pi^4 - 2\pi[\frac{y^3}{3} + \varepsilon(y\cos y - \sin y)]|_0^{2\pi} \\ &= \frac{8}{3}\pi^4 + 4\varepsilon\pi^2 \end{aligned}$$

Example 2 To calculate the volume of revolution of curvilinear trapezoid: $y = f(x) = \frac{x}{\pi} + \sin x$ $(0 \le x \le 2\pi)$, $y = 0$, $y = 2\pi$ and y-axis rotating with y-axis.



Figure 5. Kepler equation $y = x + \varepsilon \sin y$, $\varepsilon \in (0, 1)$.



Figure 6. Piecewise monotone intervals of $f(x) = \frac{x}{\pi} + \sin x, 0 \le x \le 2\pi$.

Solution 1: We give the classical solution.

(1) Let $f'(x) = (\frac{x}{\pi} + \sin x)' = \frac{1}{\pi} + \cos x = 0$, we obtain two extreme value points, $x_1 = \arccos(-\frac{1}{\pi})$ and $x_2 = 2\pi - \arccos(-\frac{1}{\pi})$. And,

$$f(x_1) = \frac{1}{\pi}\arccos(-\frac{1}{\pi}) + \sin(\arccos(-\frac{1}{\pi})),$$

$$f(x_2) = \frac{1}{\pi}[2\pi - \arccos(-\frac{1}{\pi})] - \sin(\arccos(-\frac{1}{\pi})).$$

(2) Denote:

$$\begin{array}{ll} s_1 : y = f(x), 0 \le x \le x_0. & s_2 : y = f(x), x_0 \le x \le x_1. \\ s_1 + s_2 : y = f(x), 0 \le x \le x_1. & \\ s_3 : y = f(x), x_1 \le x \le x_2. & s_4 : y = f(x), x_2 \le x \le x_3. \end{array}$$

The volumes of curved trapezoids of $s_1, s_2, s_3, s_4, s_1 + s_2$ in term of y-axis rotating with y-axis are denoted as V_{s_1},V_{s_2},V_{s_3} ,V_{s_4},$V_{s_1+s_2}$,and the volume of curved trapezoid $y = f(x)$ $(0 \le x \le 2\pi)$, $y = 0$, $y = 2$, $x = 0$ rotating with y-axis is V_Y.Then,

$$\begin{aligned} V_Y &= V_{s_4} - (V_{s_3} - V_{s_2}) + V_{s_1} = V_{s_4} - V_{s_3} + (V_{s_2} + V_{s_1}) \\ &= V_{s_4} - V_{s_3} + V_{s_1+s_2} \end{aligned} \tag{12}$$

According to Theorem 1,

$$\begin{aligned} V_{s_1+s_2} &= \pi x_1^2 f(x_1) - 0 - 2\pi \int_0^{x_1} x(\frac{x}{\pi} + \sin x)dx \\ &= \pi x_1^2 f(x_1) - 2\pi[\frac{x^3}{3\pi} + (\sin x - x\cos x)]|_0^{x_1} \\ &= \pi x_1^2 f(x_1) - 2\pi[\frac{x_1^3}{3\pi} + (\sin x_1 - x_1\cos x_1)] \end{aligned}$$

$$\begin{aligned} V_{s_3} &= 2\pi \int_{x_1}^{x_2} x(\frac{x}{\pi} + \sin x)dx - \pi[x_2^2 f(x_2) - x_1^2 f(x_1)] \\ &= 2\pi[\frac{x^3}{3\pi} + (\sin x - x\cos x)]|_{x_1}^{x_2} - \pi[x_2^2 f(x_2) - x_1^2 f(x_1)] \\ &= 2\pi[\frac{x_2^3}{3\pi} + (\sin x_2 - x_2\cos x_2)] - 2\pi[\frac{x_1^3}{3\pi} + (\sin x_1 - x_1\cos x_1)] - \pi[x_2^2 f(x_2) - x_1^2 f(x_1)] \end{aligned}$$

$$\begin{aligned} V_{s_4} &= \pi[x_3^2 f(x_3) - x_2^2 f(x_2)] - 2\pi \int_{x_2}^{x_3} x(\frac{x}{\pi} + \sin x)dx \\ &= \pi[x_3^2 f(x_3) - x_2^2 f(x_2)] - 2\pi[\frac{x^3}{3\pi} + (\sin x - x\cos x)]|_{x_2}^{x_3} \\ &= \pi[x_3^2 f(x_3) - x_2^2 f(x_2)] - 2\pi[\frac{x_3^3}{3\pi} + (\sin x_3 - x_3\cos x_3)] + 2\pi[\frac{x_2^3}{3\pi} + (\sin x_2 - x_2\cos x_2)] \end{aligned}$$

Substituting $V_{s_1+s_2}$,V_{s_3},V_{s_4},$f(x_3) = f(2\pi) = 2$ into formula (12), we obtain

$$\begin{aligned} V_Y &= V_{s_4} - V_{s_3} + V_{s_1+s_2} = \pi x_3^2 f(x_3) - 2\pi[\frac{x_3^3}{3\pi} + (\sin x_3 - x_3\cos x_3)] \\ &= 8\pi^3 - 2\pi[\frac{8\pi^2}{3} - 2\pi] = \frac{8\pi^3}{3} + 4\pi^2 \end{aligned}$$

Solution 2: Using formula (8), the volume of solids of revolution is

$$\begin{aligned} V_Y &= \text{sgn}(f(b) - f(a)) \left\{ \pi \left[b^2 f(b) - a^2 f(a) \right] - 2\pi \int_a^b x f(x)dx \right\} \\ &= \left\{ \pi \left[(2\pi)^2 f(2\pi) - 0 \right] - 2\pi \int_0^{2\pi} x(\frac{x}{\pi} + \sin x)dx \right\} \\ &= 8\pi^3 - 2\pi[\frac{x^3}{\pi} + \sin x - x\cos x]|_0^{2\pi} \\ &= 8\pi^3 - 2\pi[\frac{8\pi^2}{3} - 2\pi] = \frac{8\pi^3}{3} + 4\pi^2 \end{aligned}$$

Similar to Example 2, interchanging x and y, we can obtain an example rotating with x-axis using Theorem 3.

Example 3 To calculate the volume of curved trapezoid of $x = g(y) = \frac{y}{\pi} + \sin y$ ($0 \le y \le 2\pi$), $x = 0, x = 2, y = 0$ rotating with x-axis.

(The solution is similar to Example 2, we omit the detail.)

ACKNOWLEDGMENT. This paper is partially supported by 863 Project of China (2008AA02Z306), and Major Program of the National Natural Science Foundation of China (No.61327807).

REFERENCES

1. J. Stewart. *Calculus* (seventh edition). Brooks/Cole, Cengage Learning. 2012. 343-494.
2. V. A. Zorich. *Mathematical Analysis I* (4th edition). Springer-Verlag Berlin Heidelberg. 2004. 374-385.
3. A. Hall. Kepler's Problem. *Annals of Mathematics.***10** (3) (1883) 65–66.